

8. (a) An urn contains four tickets marked with numbers 112, 121, 211, 222 and one ticket is drawn at random. Let $A_i (i = 1, 2, 3)$ be the event that i th digit of the number of the ticket drawn is 1. Discuss the independence of the event A_1, A_2 and A_3 .
- (b) State Baye's theorem. A and B are two weak students of statistics and their chances of solving a problem in statistics correctly are $\frac{1}{6}$ and $\frac{1}{8}$ respectively. If the probability of their making a common error is $\frac{1}{525}$ and they obtain the same answer, find the probability that their answer is correct.

6, 6

This question paper contains 8 printed pages]

4/12/17
(M)

Roll No.

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

S. No. of Question Paper : 6711

Unique Paper Code : 323271101

HC

Name of the Paper : Descriptive Statistics

Name of the Course : B.Sc. (H) Statistics

Semester : I

Duration : 3 Hours

Maximum Marks : 75

(Write your Roll No. on the top immediately on receipt of this question paper.)

Attempt 6 questions in all.

Question No. 1 is compulsory.

Attempt 5 more questions selecting 3 questions from Section A

and 2 questions from Section B.

Use of simple calculator is allowed.

Section A

1. (a) Fill in the blanks :

(i) For a symmetric distribution, $\beta_1 = \dots\dots\dots$

(ii) If the attributes of A and B are independent, then $\frac{(AB)}{N} = \dots\dots\dots$

(iii) The algebraic sum of the deviations of 20 observations measured from 30 is 2. Therefore, mean of these observations is $\dots\dots\dots$

(iv) Correlation coefficient is the $\dots\dots\dots$ of the regression coefficients.

(v) Total number of ultimate class frequency for n attributes is $\dots\dots\dots$

(vi) If A and B are mutually disjoint events, then $P(A \cup B) = \dots\dots\dots$

(b) In a random arrangement of the letters of the word COMMERCE, find the probability that all the vowels come together. 6, 6

7. (a) Show that in a population with three attributes A, B and C if :

(i) $(AB) = (A)$ and $(BC) = (B)$, then $(AC) = (A)$;

(ii) If $(AB) = (A)$ and $(BC) = 0$, then $(AC) = 0$.

(b) One bag contains 5 white and 4 black balls. Another bag contains 7 white and 9 black balls. A ball is transferred from the first bag to the second bag and then a ball is drawn from the second. Find the probability that the ball is white. 6, 6

(b) If d_i be the difference in the ranks of the i th individual in two different characteristics, then show that the maximum value of :

$$\sum_{i=1}^n d_i^2 \text{ is } \frac{1}{3}(n^3 - n). \quad 6, 6$$

5. (a) If X and Y are independent random variables with zero means and standard deviations 3 and 4 respectively, then find K so that $X + 2Y$ and $KX - Y$ are uncorrelated.

(b) If X and Y are two independent variables, show that :

$$r(X + Y, X - Y) = r^2(X, X + Y) - r^2(Y, X + Y)$$

where $r(X + Y, X - Y)$ denotes the coefficient of correlation between $X + Y$ and $X - Y$. 6, 6

Section B

6. (a) State and prove Boole's inequality.

(vii). A, B and C are three mutually exclusive and exhaustive events associated with a random experiment. Given that :

$$P(B) = \frac{3}{2}P(A) \text{ and } P(C) = \frac{1}{2}P(B), \text{ then}$$

$P(B) = \dots\dots\dots$

(viii) The limits for rank correlation coefficient are

(ix) Two uncorrelated variables be independent.

(b) (i) If the two regression lines are $3x + 12y = 19$, $3y + 9x = 46$, then find correlation coefficient between x and y .

(ii) If $P(A \cup B) = \frac{5}{6}$, $P(A \cap B) = \frac{1}{3}$ and $P(\bar{A}) = \frac{1}{2}$, then find $P(A)$ and $P(B)$. Hence show that A and B are independent.

(iii) If the lines of regression of Y on X and X on Y are respectively $a_1x + b_1y + c_1 = 0$ and $a_2x + b_2y + c_2 = 0$, then prove that

$$a_1b_2 \leq a_2b_1. \quad 9 \times 1, 3 \times 2$$

2. (a) Show that the sum of the squares of the deviations of a set of values is minimum when taken about mean.

(b) In a frequency table, the upper boundary of each class interval has a constant ratio to the lower boundary.

Show that the geometric mean G may be expressed by the formula :

$$\log G = x_0 + \frac{C}{N} \sum_i f_i (i - 1).$$

Where x_0 is the logarithm of the mid-value of the first interval and C is the logarithm of the ratio between upper and lower boundaries.

6, 6

3. (a) What do you mean by skewness and kurtosis of a distribution? Explain the methods of measuring skewness and kurtosis of a frequency distribution.

(b) Let X/f_i , $i = 1, 2, 3, \dots, n$ be a discrete series. If the deviations $x_i = X_i - M$ are small compared with mean M, so that $\left(\frac{x}{M}\right)^3$ and high powers of $\left(\frac{x}{M}\right)$ are neglected,

then :

$$(i) \quad G = M \left(1 - \frac{\sigma^2}{2M^2} \right),$$

$$(ii) \quad \text{Mean}(\sqrt{x}) = \sqrt{M \left(1 - \frac{\sigma^2}{8M^2} \right)},$$

where G is the geometric mean, M is the arithmetic mean and σ is the standard deviation. 6, 6

4. (a) Explain the principle of least squares. Obtain the equation of the line of regression of Y on X.