

(c) Explain why  $f(x) = \sin(x)$  is a bounded function on  $\mathbf{R}$ , whereas  $f(z) = \sin(z)$  is not a bounded function on the complex plane  $\mathbf{C}$ , although  $\sin^2(z) + \cos^2(z) = 1$  for all  $z \in \mathbf{C}$ .

4. (a) State and prove Cauchy Integral formula.

(b) State Liouville's theorem and use it to prove the fundamental theorem of algebra.

(c) Let  $C$  denote a contour of length  $L$ , and suppose that a function  $f(z)$  is piecewise continuous on  $C$ . Show that

$$\left| \int_C f(z) dz \right| \leq ML, \text{ where } M \text{ is a non-negative constant}$$

such that  $|f(z)| \leq M$  for all points  $z$  on  $C$  at which  $f(z)$  is defined. Hence, show that

$$\left| \int_{C_R} \frac{2z^2 - 1}{z^4 + 5z^2 + 4} dz \right| \leq \frac{\pi R(2R^2 + 1)}{(R^2 - 1)(R^2 - 4)}, \text{ where } C_R$$

denote the upper half of the circle  $|z| = R (R > 2)$ , taken in the counterclockwise direction.

5. (a) Derive the expansions :

$$(i) \frac{\sinh z}{z^2} = \frac{1}{z} + \sum_{n=0}^{\infty} \frac{z^{2n+1}}{(2n+3)!} \quad (0 < |z| < \infty);$$

$$(ii) z^3 \cosh\left(\frac{1}{z}\right) = \frac{z}{2} + z^3 + \sum_{n=1}^{\infty} \frac{1}{(2n+2)!}$$

$$\frac{1}{z^{2n-1}} \quad (0 < |z| < \infty)$$

Friday 10/5/2019

This question paper contains 4+2 printed pages]

Roll No.

S. No. of Question Paper : 2253

Unique Paper Code : 32351601

IC

Name of the Paper : Complex Analysis

Name of the Course : B.Sc. (Hons.) Mathematics

Semester : VI

Duration : 3 Hours

Maximum Marks : 75

(Write your Roll No. on the top immediately on receipt of this question paper.)

Attempt all parts from Question No. 1.

Each part carries 1½ marks.

Attempt any two parts from question Nos. 2 to 6

Each part carries six marks.

1. State True or False. Justify your answer in brief :

(a) A point  $z_0$  of a domain need not be an accumulation point of that domain.

(b)  $\lim_{z \rightarrow 0} \frac{\bar{z}^2}{z} = 0$

P.T.O.

- (c) The function  $f(z) = e^z$  is periodic with period  $2\pi$ .
- (d)  $\log(-ei) = 1 - \frac{\pi}{2}i$ .
- (e) The function  $f(z) = |z|^2$  is analytic at  $z = 0$ .
- (f) Let  $C$  denote the boundary of the triangle with vertices at the point  $0$ ,  $3i$ , and  $-4$ , oriented in the counterclockwise direction. Then  $|\int_C (e^z - \bar{z}) dz| \leq 60$ .
- (g) If  $C$  is any simple closed contour, in either direction, then  $\int_C \exp(z^3) dz = 0$ .
- (h) If  $C$  is the positively oriented unit circle  $|z| = 1$ , then  $\int_C \frac{\exp(2z)}{z^4} dz = \frac{8\pi i}{3}$ .
- (i)  $\text{Res}_{z=0} f(z) = -\frac{1}{3!}$ , where  $f(z) = z^2 \sin\left(\frac{1}{z}\right)$ ,  $0 < |z| < \infty$ .
- (j) The function  $f(z) = \frac{1}{\sin\left(\frac{\pi}{z}\right)}$  has no isolated singular point.

2. (a) Prove that a finite set of points cannot have any accumulation point.
- (b) Suppose that  $f(z) = u(x, y) + iv(x, y)$  ( $z = x + iy$ ) and  $z_0 = x_0 + iy_0$ ,  $w_0 = u_0 + iv_0$ . Then  $\lim_{z \rightarrow z_0} f(z) = w_0$  if and only if  $\lim_{(x,y) \rightarrow (x_0,y_0)} u(x, y) = u_0$  and  $\lim_{(x,y) \rightarrow (x_0,y_0)} v(x, y) = v_0$ .
- (c) Define neighbourhood of the point at infinity. Show that a set  $S$  is unbounded if and only if every neighbourhood of the point at infinity contains at least one point in  $S$ .
3. (a) Use Cauchy-Riemann equations to show that  $f'(z)$  does not exist at any point if  $f(z) = \exp(\bar{z})$ . State sufficient conditions for differentiability of a function  $f(z)$  at any point  $z_0 = x_0 + iy_0 \in C$ .
- (b) Suppose that a function  $f(z) = u(x, y) + iv(x, y)$  and its conjugate  $\overline{f(z)} = u(x, y) - iv(x, y)$  are both analytic in a given domain  $D$ . Show that  $f(z)$  must be constant throughout  $D$ .