-02112/2019(M6.) 2/12/2019(M)

[This question paper contains 7 printed pages]

Your Roll No. Sl. No. of Q. Paper

parties fruct Y is risson.

the states and the states

Unique Paper Code

Name of the Course

Name of the Paper

: Metric Spaces : V

: 7466

: 32351501

: B.Sc.(Hons.) **Mathematics**

Semester

Alter mitt at the distance in the task

Time : 3 Hours

Maximum Marks: 75

J

are at the X **Instructions for Candidates :**

- (a) Write your Roll No. on the top immediately on receipt of this question paper.
- (b) Attempt any two parts from each question.

1. (a) Define a metric space. Let $p \ge 1$. Define $d_p: \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R} \text{ as } d_p(x, y) = \left(\sum_{i=1}^n |x_i - y_i|^p\right)^{1/p}$, $x (x_1, x_2, ..., x_n), y = (y_1, y_2, ..., y_n) \in \mathbb{R}^n$. Show that (\mathbb{R}^n, d_p) is a metric space.

6.5

P.T.O.

na ploele

- (b) When is a metric space said to be complete ? Is discrete metric space complete ? Justify. 6.5
- (c) Let (X, d) be a metric space. Define $d_1: X \times X$

 $\rightarrow \mathbb{R} \text{ by } d_1(x,y) = \frac{d(x,y)}{1+d(x,y)}, \text{ for all } x, y \in X.$ Prove that d_1 is a metric on X and d_1 is equivalent to d.

2. (a) Prove that every open ball in a metric space

(X, d) is an open set in (X, d). What aboutthe converse ? Justify.

(b) Define a homeomorphism from a metric

space (X,d_1) to a metric space (Y, d_2) . Show that the function $f: \mathbb{R} \to]-1$, 1[defined by

6

 $f(x) = \frac{x}{1+|x|}$ is a homeomorphism.

2

7466

3500

- (b) Let (X, d) be a metric space and Y be a subset of X. If Y is compact subset of (X, d), then prove that Y is closed.
- (c) Let f be a continuous function from a compact metric space (X,d₁) to a metric space (Y, d₂). Prove that f is uniformly continuous on X.

- (c) Let (X, d) be a metric space. Then prove that following statements are equivalent : $1.5 \times 4 = 6$
 - (X, d) is disconnected. (i)
 - There exist two non-empty disjoint (ii) subsets A and B, both open in X, such that $X = A \bigcup B$.
 - There exist two non-empty disjoint (iii) subsets A and B, both closed in X, such that X = A[]B.
 - There exists a proper subset of X, which (iv) is both open and closed in X.

6.5

6. (a) Let (\mathbb{R}, d) be the space of real numbers with the usual metric. Show that a connected subset of \mathbb{R} must be an interval. Give example of two connected subsets of $\mathbb R$ such that their union is disconnected.

- (c) Let (X, d) be a metric space and let A, B be non-empty subsets of X. Prove that : 6
 - (i) $(A \cap B)^0 = A^0 \cap B^0$ (ii) $\overline{A \cup B} = \overline{A} \cup \overline{B}$
- (a) Let (X, d) be a metric space and $F \subseteq X$. Prove 3. that the following statements are equivalent :
 - (i) $x \in \overline{F}$
 - (ii) $S(x,\varepsilon) \cap F \neq \phi$, for every open ball $S(x,\varepsilon)$ centred at x
 - (iii) There exists an infinite sequence $\{x_n\}$ of point (not necessarily distinct) of F such that $X_n \rightarrow X.$
 - (b) Let (X,d) be a metric space and $F \subseteq X$. Prove that F is closed in X if and only if F^c is open in X. where F^c is complement of F in X.

3

6

P.T.O.

- (c) Let (X,d) be a metric space such that for every nested sequence {F_n}_{n≥1} of non-empty closed subsets of X satisfying d(F_n) → 0 as n →∞, the intersection ∩[∞]_{n=1}F_n contains exactly one point. Prove that (X,d) is complete.
- 4. (a) Let f be a mapping from a metric space (X, d₁) to a metric space (Y, d₂). Prove that f is continuous on X if and only if f⁻¹ (G) is open in X for all open subsets G of Y. 6.5
 - (b) Let (X, d₁) and (Y, d₂) be two metric spaces.
 Prove that the following statements are equivalent : 6.5
 - (i) f is continuous on X
 - (ii) $\overline{f^{-1}(B)} \subseteq \overline{f^{-1}(\overline{B})}$, for all subsets B of Y (iii) $f(\overline{A}) \subseteq \overline{f(A)}$, for all subsets A of X.

- (c) Define uniform continuity of a function f from a metric space (X, d₁) to a metric space (Y, d₂). Let (X, d) be a metric space and A be a non-empty subset of X. Show that the function f: (X, d) → R defined as f (x) = d (x, A), for all x∈X, is uniformly continuous on X.
 - 6.5

- 5. (a) State and prove contraction mapping theorem.
 - (b) (i) Let Y be a non-empty subset of a metric space (X, d) and (Y,d_y) be complete, where d_y is restriction of d to Y × Y. Prove that Y is closed in X.
 (ii) Let A be a non-empty bounded subset of a metric space (X, d). Prove that

- $d(A) = d(\overline{A}).$
 - P.T.O.

-021.12/19 (Mosn.) 2/12/2019(m)

•

[This question paper contains 7 printed pages]

: 7466

: 32351501

: B.Sc.(Hons.) **Mathematics**

: Metric Spaces

1290 ne m (b . X

Your Roll No.

prove that Y is closed.

with he a promotional for the filling

Sl. No. of Q. Paper

Unique Paper Code

Name of the Course

Name of the Paper

Semester

12110 000

Time : 3 Hours

Maximum Marks: 75

J

Instructions for Candidates :

1830

(a) Write your Roll No. on the top immediately on receipt of this question paper.

: V

(b) Attempt any two parts from each question.

1. (a) Define a metric space. Let $p \ge 1$. Define $d_p: \mathbb{R}^n \times \mathbb{R}^n \to \mathbb{R} \text{ as } d_p(x, y) = \left(\sum_{i=1}^n |x_i - y_i|^p\right)^{1/p},$ $x (x_1, x_2, ..., x_n), y = (y_1, y_2, ..., y_n) \in \mathbb{R}^n$. Show that (\mathbb{R}^n, d_p) is a metric space.

6.5

P.T.O.

- (b) When is a metric space said to be complete ? Is discrete metric space complete ? Justify. 6.5
- (c) Let (X, d) be a metric space. Define $d_1: X \times X$

 $\rightarrow \mathbb{R} \text{ by } d_1(x,y) = \frac{d(x,y)}{1+d(x,y)}, \text{ for all } x, y \in X.$ Prove that d_1 is a metric on X and d_1 is equivalent to d_1 .

- 2. (a) Prove that every open ball in a metric space
 (X, d) is an open set in (X, d). What about the converse ? Justify.
 - (b) Define a homeomorphism from a metric space (X,d_1) to a metric space (Y, d_2) . Show that the function $f: \mathbb{R} \to]-1$, 1[defined by

 $f(x) = \frac{x}{1+|x|}$ is a homeomorphism.

- (b) Let (X, d) be a metric space and Y be a subset of X. If Y is compact subset of (X, d), then prove that Y is closed.
- (c) Let f be a continuous function from a compact metric space (X,d₁) to a metric space (Y, d₂). Prove that f is uniformly continuous on X.

7

6.5

- (c) Let (X, d) be a metric space. Then prove that following statements are equivalent : 1.5×4=6
 - (i) (X, d) is disconnected.
 - (ii) There exist two non-empty disjoint subsets A and B, both open in X, such that X = A∪B.
 - (iii) There exist two non-empty disjoint subsets A and B, both closed in X, such that X = A∪B.
 - (iv) There exists a proper subset of X, which is both open and closed in X.
- 6. (a) Let (R, d) be the space of real numbers with the usual metric. Show that a connected subset of R must be an interval. Give example of two connected subsets of R such that their union is disconnected.

- (c) Let (X, d) be a metric space and let A, B be non-empty subsets of X. Prove that : 6
 (i) (A∩B)⁰ = A⁰∩B⁰
 - (ii) $\overline{A \cup B} = \overline{A} \cup \overline{B}$
- (a) Let (X, d) be a metric space and F⊆X. Prove that the following statements are equivalent :
 - 6

(i) $x \in \overline{F}$

- (ii) S(x,ε)∩F≠φ, for every open ball S(x,ε) centred at x
- (iii) There exists an infinite sequence $\{x_n\}$ of point (not necessarily distinct) of F such that $x_n \rightarrow X$.
- (b) Let (X,d) be a metric space and F⊆X. Prove that F is closed in X if and only if F^c is open in X, where F^c is complement of F in X.

3

6.5

P.T.O.

(c) Let (X,d) be a metric space such that for every nested sequence {F_n}_{n≥1} of non-empty closed subsets of X satisfying d(F_n) → 0 as n →∞, the intersection ∩[∞]_{n=1}F_n contains exactly one point. Prove that (X,d) is complete.

(

- 4. (a) Let f be a mapping from a metric space (X, d₁) to a metric space (Y, d₂). Prove that f is continuous on X if and only if f⁻¹ (G) is open in X for all open subsets G of Y. 6.5
 - (b) Let (X, d₁) and (Y, d₂) be two metric spaces.
 Prove that the following statements are equivalent : 6.5
 - (i) f is continuous on X
 - (ii) $\overline{f^{-1}(B)} \subseteq \overline{f^{-1}(\overline{B})}$, for all subsets B of Y (iii) $f(\overline{A}) \subseteq \overline{f(A)}$, for all subsets A of X.

(c) Define uniform continuity of a function f from a metric space (X, d₁) to a metric space (Y, d₂). Let (X, d) be a metric space and A be a non-empty subset of X. Show that the function f: (X, d) → R defined as f (x) = d (x, A), for all x∈X, is uniformly continuous on X.

6.5

5. (a) State and prove contraction mapping theorem.
(b) (i) Let Y be a non-empty subset of a metric space (X, d) and (Y,d_y) be complete, where d_y is restriction of d to Y × Y. Prove that Y is closed in X.
(ii) Let A be a non-empty bounded subset of a metric space (X, d). Prove that d(A) = d(A

5 P.T.O.