


Areas Under the One-Tailed Standard Normal Curve

This table provides the area between the mean and some Z score. For example, with a Z score = 1.45 the area = 0.4265.



Z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.0000	0.0040	0.0080	0.0120	0.0160	0.0199	0.0239	0.0279	0.0319	0.0359
0.1	0.0398	0.0438	0.0478	0.0517	0.0557	0.0596	0.0636	0.0675	0.0714	0.0753
0.2	0.0793	0.0832	0.0871	0.0910	0.0948	0.0987	0.1026	0.1064	0.1103	0.1141
0.3	0.1179	0.1217	0.1255	0.1293	0.1331	0.1368	0.1406	0.1443	0.1480	0.1517
0.4	0.1554	0.1591	0.1628	0.1664	0.1700	0.1736	0.1772	0.1808	0.1844	0.1879
0.5	0.1915	0.1950	0.1985	0.2019	0.2054	0.2088	0.2123	0.2157	0.2190	0.2224
0.6	0.2257	0.2291	0.2324	0.2357	0.2389	0.2422	0.2454	0.2486	0.2517	0.2549
0.7	0.2580	0.2611	0.2642	0.2673	0.2704	0.2734	0.2764	0.2794	0.2823	0.2852
0.8	0.2881	0.2910	0.2939	0.2967	0.2995	0.3023	0.3051	0.3078	0.3106	0.3133
0.9	0.3159	0.3187	0.3212	0.3238	0.3264	0.3289	0.3315	0.3340	0.3365	0.3389
1.0	0.3413	0.3438	0.3461	0.3485	0.3508	0.3532	0.3554	0.3577	0.3599	0.3621
1.1	0.3643	0.3665	0.3686	0.3708	0.3729	0.3749	0.3770	0.3790	0.3810	0.3830
1.2	0.3849	0.3869	0.3888	0.3907	0.3925	0.3944	0.3962	0.3980	0.3997	0.4015
1.3	0.4032	0.4049	0.4066	0.4082	0.4099	0.4115	0.4131	0.4147	0.4162	0.4177
1.4	0.4192	0.4207	0.4222	0.4236	0.4251	0.4265	0.4279	0.4292	0.4306	0.4319
1.5	0.4332	0.4345	0.4357	0.4370	0.4382	0.4394	0.4406	0.4418	0.4429	0.4441
1.6	0.4452	0.4463	0.4474	0.4484	0.4495	0.4505	0.4515	0.4525	0.4535	0.4545
1.7	0.4554	0.4564	0.4573	0.4582	0.4591	0.4599	0.4608	0.4616	0.4625	0.4633
1.8	0.4641	0.4649	0.4656	0.4664	0.4671	0.4678	0.4686	0.4693	0.4699	0.4705
1.9	0.4713	0.4719	0.4726	0.4732	0.4738	0.4744	0.4750	0.4756	0.4761	0.4767
2.0	0.4772	0.4778	0.4783	0.4788	0.4793	0.4798	0.4803	0.4808	0.4812	0.4817
2.1	0.4821	0.4826	0.4830	0.4834	0.4838	0.4842	0.4846	0.4850	0.4854	0.4857
2.2	0.4861	0.4864	0.4868	0.4871	0.4875	0.4878	0.4881	0.4884	0.4887	0.4890
2.3	0.4893	0.4896	0.4898	0.4901	0.4904	0.4906	0.4909	0.4911	0.4913	0.4916
2.4	0.4918	0.4920	0.4922	0.4925	0.4927	0.4929	0.4931	0.4932	0.4934	0.4936
2.5	0.4938	0.4940	0.4941	0.4943	0.4945	0.4946	0.4948	0.4949	0.4951	0.4952
2.6	0.4953	0.4955	0.4956	0.4957	0.4959	0.4960	0.4961	0.4962	0.4963	0.4964
2.7	0.4965	0.4966	0.4967	0.4968	0.4969	0.4970	0.4971	0.4972	0.4973	0.4974
2.8	0.4974	0.4975	0.4976	0.4977	0.4977	0.4978	0.4979	0.4979	0.4980	0.4981
2.9	0.4981	0.4982	0.4982	0.4983	0.4984	0.4984	0.4985	0.4985	0.4986	0.4986
3.0	0.4987	0.4987	0.4987	0.4988	0.4988	0.4989	0.4989	0.4989	0.4990	0.4990
3.1	0.4990	0.4991	0.4991	0.4991	0.4992	0.4992	0.4992	0.4992	0.4992	0.4993
3.2	0.4993	0.4993	0.4994	0.4994	0.4994	0.4994	0.4994	0.4995	0.4995	0.4995
3.3	0.4995	0.4995	0.4995	0.4996	0.4996	0.4996	0.4996	0.4996	0.4996	0.4997
3.4	0.4997	0.4997	0.4997	0.4997	0.4997	0.4997	0.4997	0.4997	0.4997	0.4998
3.5	0.4998	0.4998	0.4998	0.4998	0.4998	0.4998	0.4998	0.4998	0.4998	0.4998
3.6	0.4998	0.4998	0.4999	0.4999	0.4999	0.4999	0.4999	0.4999	0.4999	0.4999
3.7	0.4999	0.4999	0.4999	0.4999	0.4999	0.4999	0.4999	0.4999	0.4999	0.4999
3.8	0.4999	0.4999	0.4999	0.4999	0.4999	0.4999	0.4999	0.4999	0.4999	0.4999
3.9	0.5000	0.5000	0.5000	0.5000	0.5000	0.5000	0.5000	0.5000	0.5000	0.5000

[This question paper contains 8 printed pages.]

Your Roll No.....

Sr. No. of Question Paper : 3034

D

Unique Paper Code : 2922061102

Name of the Paper : Statistics for Business Decision

Name of the Course : BMS

Semester : I

Duration : 3 Hours

Maximum Marks : 90

Instructions for Candidates

- Write your Roll No. on the top immediately on receipt of this question paper.
- All questions carry equal marks.
- Use of Simple calculators are allowed.
- Reference to Probability distribution tables (Binomial, Poisson and Normal) and Student T-test table is allowed.
- (a) Given below is the data of daily income of workers employed in a manufacturing unit:

Daily Income of Workers (in Rs)	No of Workers
100-150	4
150-200	10
200-250	15
250-300	25
300-350	8
350-400	18

Compute (a) highest income of lowest 50% workers (b) Modal wages and (c) minimum income of top 25% workers. (9)

(b) Enumerate the properties of regression coefficients. Given the regression equation of Y on X and X on Y are respectively $Y=2X$ and $6X-Y=4$ and the second moment of X about the origin is 3. Find the correlation coefficient and standard deviation of Y. (9)

2. (a) List the conditions under which a binomial distribution can be approximated as a Poisson distribution. A production process is expected to generate 5 defectives in a batch of 100 units. Find the mean number of defectives and variance. Estimate the probability that there will be (i) no defectives, (ii) at most 2 defectives? (9)

t Table

cum. prob	$t_{.50}$	$t_{.25}$	$t_{.20}$	$t_{.15}$	$t_{.10}$	$t_{.05}$	$t_{.025}$	$t_{.01}$	$t_{.005}$	$t_{.001}$	$t_{.0005}$
	one-tail	two-tails	two-tails	two-tails	two-tails	two-tails	two-tails	two-tails	two-tails	two-tails	two-tails
df	0.50	0.25	0.20	0.15	0.10	0.05	0.025	0.01	0.005	0.001	0.0005
1	0.000	1.000	1.375	1.963	3.078	6.314	12.71	31.82	63.66	318.31	636.62
2	0.000	0.816	1.061	1.386	1.886	2.920	4.303	6.985	9.825	22.327	31.999
3	0.000	0.765	0.978	1.250	1.638	2.353	3.182	4.541	5.841	10.215	12.924
4	0.000	0.741	0.941	1.190	1.533	2.132	2.776	3.747	4.604	7.173	8.610
5	0.000	0.727	0.920	1.156	1.476	2.015	2.571	3.365	4.032	5.893	6.869
6	0.000	0.718	0.906	1.134	1.440	1.943	2.447	3.143	3.707	5.208	5.858
7	0.000	0.711	0.896	1.119	1.415	1.895	2.365	2.998	3.499	4.785	5.408
8	0.000	0.706	0.889	1.108	1.397	1.860	2.306	2.896	3.355	4.501	5.041
9	0.000	0.703	0.883	1.100	1.383	1.833	2.262	2.821	3.250	4.297	4.781
10	0.000	0.700	0.879	1.093	1.372	1.812	2.228	2.784	3.169	4.144	4.587
11	0.000	0.697	0.876	1.088	1.363	1.796	2.201	2.718	3.100	4.025	4.437
12	0.000	0.695	0.873	1.083	1.356	1.782	2.179	2.681	3.055	3.930	4.318
13	0.000	0.694	0.870	1.079	1.350	1.771	2.160	2.650	3.012	3.852	4.221
14	0.000	0.692	0.868	1.076	1.345	1.761	2.145	2.624	2.977	3.787	4.140
15	0.000	0.691	0.866	1.074	1.341	1.753	2.131	2.602	2.947	3.733	4.073
16	0.000	0.690	0.865	1.071	1.337	1.746	2.120	2.583	2.921	3.686	4.015
17	0.000	0.689	0.863	1.069	1.333	1.740	2.110	2.567	2.896	3.646	3.965
18	0.000	0.688	0.862	1.067	1.330	1.734	2.101	2.552	2.878	3.610	3.922
19	0.000	0.688	0.861	1.066	1.328	1.729	2.093	2.539	2.861	3.579	3.883
20	0.000	0.687	0.860	1.064	1.325	1.725	2.086	2.528	2.845	3.552	3.850
21	0.000	0.686	0.859	1.063	1.323	1.721	2.080	2.518	2.831	3.527	3.819
22	0.000	0.686	0.858	1.061	1.321	1.717	2.074	2.508	2.819	3.505	3.792
23	0.000	0.685	0.858	1.060	1.319	1.714	2.069	2.500	2.807	3.485	3.766
24	0.000	0.685	0.857	1.059	1.318	1.711	2.064	2.492	2.797	3.467	3.745
25	0.000	0.684	0.856	1.058	1.316	1.708	2.060	2.485	2.787	3.450	3.725
26	0.000	0.684	0.856	1.058	1.315	1.706	2.056	2.479	2.779	3.435	3.707
27	0.000	0.684	0.855	1.057	1.314	1.703	2.052	2.473	2.771	3.421	3.690
28	0.000	0.683	0.855	1.056	1.313	1.701	2.048	2.467	2.763	3.408	3.674
29	0.000	0.683	0.854	1.055	1.311	1.699	2.045	2.462	2.756	3.396	3.659
30	0.000	0.683	0.854	1.055	1.310	1.697	2.042	2.457	2.750	3.385	3.646
40	0.000	0.681	0.851	1.050	1.303	1.684	2.021	2.423	2.704	3.307	3.551
60	0.000	0.679	0.848	1.045	1.296	1.671	2.000	2.390	2.660	3.232	3.460
80	0.000	0.678	0.846	1.043	1.292	1.664	1.990	2.374	2.639	3.195	3.416
100	0.000	0.677	0.845	1.042	1.290	1.660	1.984	2.364	2.626	3.174	3.390
1000	0.000	0.675	0.842	1.037	1.282	1.646	1.962	2.330	2.581	3.098	3.300
Z	0.000	0.674	0.842	1.036	1.282	1.645	1.960	2.326	2.576	3.090	3.291
	0%	50%	60%	70%	80%	90%	95%	96%	99%	99.8%	99.9%
	Confidence Level										

(b) Distinguish between point estimation and interval estimation of population mean. Given that the income of 25 executives in an industry are approximately normally distributed, with a mean of ₹ 50,500 per month, find at 95% confidence interval if

- (i) the population standard deviation is known to be ₹ 8,900, and
 (ii) the sample standard deviation is computed as ₹ 9,500. (9)

(c) Elaborate on the properties of arithmetic mean. The first of the two samples have 100 items with mean 15 and standard deviation 3. If the whole group has 250 items with mean 15.6 and variance 13.44, find the standard deviation of the second group. (9)

Poisson Distribution Table

$\lambda =$	0.5	1.0	1.5	2.0	2.5	3.0	3.5	4.0	4.5	5.0
X=0	0.6065	0.3679	0.2231	0.1353	0.0821	0.0498	0.0302	0.0183	0.0111	0.0067
1	0.9098	0.7358	0.5578	0.4060	0.2873	0.1991	0.1359	0.0916	0.0611	0.0404
2	0.9856	0.9197	0.9197	0.8088	0.6767	0.5438	0.4232	0.3208	0.2381	0.1247
3	0.9982	0.9810	0.9344	0.8571	0.7576	0.6472	0.5366	0.4335	0.3423	0.2650
4	0.9998	0.9963	0.9814	0.9473	0.8912	0.8153	0.7254	0.6288	0.5321	0.4405
5	1.0000	0.9994	0.9994	0.9955	0.9834	0.9161	0.8576	0.7851	0.7029	0.6160
6	1.0000	0.9999	0.9991	0.9955	0.9858	0.9665	0.9347	0.8893	0.8311	0.7622
7	1.0000	1.0000	0.9998	0.9989	0.9958	0.9881	0.9733	0.9489	0.9134	0.8666
8	1.0000	1.0000	1.0000	0.9998	0.9989	0.9962	0.9901	0.9786	0.9597	0.9319
9	1.0000	1.0000	1.0000	1.0000	0.9997	0.9989	0.9967	0.9919	0.9829	0.9682
10	1.0000	1.0000	1.0000	1.0000	0.9999	0.9997	0.9990	0.9972	0.9933	0.9863

(b) Does correlation always signify a cause-and-effect relationship between the variables? Given below are scores of ten potential candidates in two different tests undertaken to measure aptitude in fine arts:

Test A	35	54	80	95	73	73	35	91	83	81
Test B	40	60	75	90	70	75	38	95	75	71

Calculate the Spearman's rank correlation for the scores and interpret the result. (9)

3. Attempt **any two** of the following :

(a) Distinguish between Qualitative and Quantitative data. Draw a Box and whisker plot for the following data pertaining to daily maximum temperature observations over a period of 20 days:

19	20	25	26	33	32	23	24	36	29
10	32	34	22	31	32	39	25	24	17

Calculate the mean and their standard deviation. Comment on the skewness of the distribution.

(9)

- (b) In a post office, three clerks are assigned to process incoming mail. The first clerk, A, processes 40 percent; the second clerk, B, processes 35 percent; and the third clerk, C, processes 25 percent of the mail. The first clerk has an error rate of 0.04, the second has an error rate of 0.06, and the third has an error rate of 0.03. A mail selected at random from a day's output is found to have an error. The postmaster wishes to know the probability that it was processed by clerk B or clerk C. What is the probability of the mail containing an error? (9)
- (c) (i) Define correlation and discuss its properties.
- (ii) The coefficient of determination of a set of data points is 0.709 and the slope of the regression line is -3.58 . Determine the correlation coefficient of the data.
- (iii) The correlation coefficient between X and Y for 20 items is 0.3. The mean of X is 15 and that of Y is 20 while the respective standard deviations are 4 and 5. At the time of calculation, one item 27 has wrongly been taken as 17 in the case of X series and 35 instead of 30 in Y series. Find the correct correlation coefficient. (9)

4. (a) Describe a normal distribution and area under the standard normal curve with the help of a diagram. The data pertaining to SAT scores was gathered to analyse the same for students in a new test preparation course. The data followed a normal distribution with a mean score of 1130 and a standard deviation of 150. Find the probability for the SAT scores exceeding 1250. (9)
- (b) Distinguish between Type I and Type II error. The breaking strengths of cables produced by a manufacturer have mean 1800 lb and standard deviation 100 lb. By a new technique in the manufacturing process, it is claimed that the breaking strength can be increased. To test this claim, a sample of 50 cables is tested, and it is found that the mean breaking strength is 1850 lb. Can we support the claim at 1% level of significance? (9)
5. Attempt **any two** of the following :
- (a) Ten items of a sample had the following values: 45, 47, 50, 52, 48, 47, 49, 53, 49, and 50. Can this sample be regarded as taken from the population having 47 as mean? (9)