This question paper contains 4 printed pages]

Roll No.		1 4			(3)	

Give examinate of function

and no manufactory therefore,

S. No. of Question Paper : 5803

Unique Paper Code : 2352011202

adjects and a Markett of the district

Name of the Paper : Calculus-DSC 5

Name of the Course : Bachelor of Science (Honours Course)

Mathematics NEP UGCF-2022

Semester : II

Duration: 3 Hours Maximum Marks: 90

(Write your Roll No. on the top immediately on receipt of this question paper.)

Attempt all questions by selecting three parts from each question.

All questions carry equal marks.

Use of calculator is not allowed.

1. (a) Using $\varepsilon - \delta$ definition of limit, show that:

$$\lim_{x \to c} \frac{1}{x} = \frac{1}{c},$$

if c > 0.

as well as differentiable on M

(b) Show that if $f:(a, \infty) \to \mathbb{R}$ (a > 0) is such that

$$\lim_{x\to\infty}xf(x)=L,$$

where $L \in \mathbf{R}$, then

$$\lim_{x\to\infty}f(x)=0.$$

Justily your answer.



- Let $A \subset \mathbf{R}$ and $f: A \to \mathbf{R}$ has a limit at $c \in \mathbf{R}$, where c is a cluster (c) point of A. Then prove that f is bounded on some neighbourhood of c.
- Give example of functions f and g such that f and g do not have limits (d)at a point c, but both f + g and fg have limits at c.
- If f is continuous on a closed interval [a, b], then prove that f is 2. (a) uniformly continuous on [a, b].
 - Show that the function $f(x) = x^2$ is uniformly continuous on [-4, 4]. **(b)**
 - Show that: (c)

$$\lim_{x\to\infty}\frac{x+2}{\sqrt{x}}(x>0)$$

does not exist in R.

Use $\varepsilon - \delta$ definition of limit to show that : (d)

$$\lim_{x \to 6} \frac{x^2 - 3x}{x + 3} = 2.$$

(a) Let 3.

$$f(x) = \begin{cases} x^2, & x \ge 0, \\ 0, & x < 0. \end{cases}$$

Calculate f' on \mathbf{R} . Is f' continuous as well as differentiable on \mathbf{R} ? Justify your answer.

Let $f(x) = \sin |x|$, where $x \in \mathbf{R}$. Determine the set of points where f (b) is not differentiable. Justify your answer.

- (c) If $f: \mathbf{R} \to \mathbf{R}$ is differentiable at $a \in \mathbf{R}$, then show that:
- (i) $\lim_{h\to 0} \frac{f(a+h)-f(a)}{h} = f'(a),$
 - (ii) $\lim_{h\to 0} \frac{f(a+h)-f(a-h)}{2h} = f'(a).$
 - (d) Suppose f is defined on an open interval (a, b) Let $x_0 \in (a, b)$. If f assumes its maximum at x_0 and if f is differentiable at x_0 , then show that $f'(x_0) = 0$.
 - 4. (a) State and prove Rolle's theorem.
 - (b) Let f be defined on **R** and suppose that $|f(x) f(y)| \le |x y|^3$ for all $x, y \in \mathbf{R}$. Prove that f is a constant function on **R**.

Truce the curve

- (c) Show that $\sin x \le x$ for all $x \ge 0$.
- (d) State Intermediate value theorem for derivatives. Suppose f is differentiable on \mathbf{R} and f(0) = 1, f(1) = 2 and f(2) = 2.
 - (i) Show that $f'(x) = \frac{1}{2}$ for some $x \in (0, 2)$,
 - (ii) Show that $f'(x) = \frac{1}{3}$ for some $x \in (0, 2)$.
 - 5. (a) If

$$y = \log\left(x + \sqrt{1 + x^2}\right),\,$$

show that

$$(1 + x^2)y_{n+2} + (2n + 1)xy_{n+1} + n^2y_n = 0.$$

(b) Trace the curve:

$$4x^2 = y^2(x^2 - 4).$$

(c) Determine the intervals of concavity and points of inflexion (if any) of the curve :

$$y = 2x^4 - 3x^2 + 2x + 1.$$

- (d) State Taylor's Theorem. Find the Taylor's series expansion of $\sin 2x$ about x = 0.
- 6. (a) Trace the curve:

$$r=2\sin 2\theta$$
. The particle is $r=2\sin 2\theta$.

(b) Determine the position and nature of the double points of the curve:

$$x^3 + 2x^2 + 2xy - y^2 + 5x - 2y = 0.$$

(c) Find the horizontal asymptotes of the graph of the function given by

$$y = x^{5} \left[\sin \frac{1}{x} - \frac{1}{x} + \frac{1}{6x^{3}} \right].$$

(d) Find the equations of tangents at origin to the curve:

$$x^4 + y^4 = a^2(x^2 - y^2).$$

